

PREPARATION OF SUBSTITUTED ALKALI METAL PIPERIDINE-4-CARBOXYLATES;
HYDROLYSIS OF 1,3,8,-TRIAZASPIRO 4.5!DECANE-2,4-DIONES, CHEMICAL
INTERMEDIATES FOR OPIUM TYPE ANALGESICS

Document Type: UTILITY

Inventors: Brackeen Marcus F (US); Feldman Paul L (US)

Assignee: Glaxo Inc Assignee Code: 25664

Preissuance Assignment Actions: Reel: 005193 Frame: 0591

Patent Number	Issue Date	Applic Number	Applic Date
------------------	---------------	------------------	----------------

Patent:	US 5039804	19910813	US 89450091	19891213
---------	------------	----------	-------------	----------

(Cited in 002 later patents)

Priority Applic:			US 89450091	19891213
------------------	--	--	-------------	----------


Calculated Expiration: 20091213

Abstract:


A method of preparing certain substituted alkali metal 4carboxy-piperidine salts (I) by alkaline hydrolysis of certain novel substituted 1,3,8-triazaspiro(4.5)decane-2, 4-diones (IV) which in turn are obtained from certain novel substituted chlorosulphonylamido piperidines (III).

Exemplary Claim:

1. The method of preparing a substituted alkali metal piperidine-4-carboxylate of the following formula:




wherein each Ar is phenyl, halophenyl, loweralkylphenyl, loweralkoxyphenyl or trifluoromethylphenyl, M is alkali metal and n is the integer 1 or 2, which comprises hydrolyzing a substituted 1,3,8-triazaspiro(4.5)decane-2, 4-dione of the following formula:




with high molar strong base having an alkali metal cationic moiety in a sealed vessel at elevated temperature to yield said substituted alkali metal piperidine-4-carboxylate.


4. The method for preparing a substituted alkali metal piperidine-4-carboxylate of the following formula (I):



wherein each Ar is phenyl, halophenyl, loweralkylphenyl, loweralkoxyphenyl or trifluoromethylphenyl, M is alkali metal and n is the integer 1 or 2 which comprises: (a) treating a substituted chlorosulphonylamido piperidine of the following formula (III):



with aqueous acid solution at elevated temperature, followed by cooling and adjustment of the pH with base, to yield the substituted 1,3,8-triaz-aspiro(4,5)decane-2,4-dione of the following formula (IV):



(b) and hydrolyzing (IV) with high molar strong base having an alkali metal cationic moiety in a sealed vessel at elevated temperature to yield the substituted alkali metal piperidine-4carboxylate of formula (I).

BEST AVAILABLE COPY